

Components improve the diagnosis of wheat allergy

Identify primary wheat sensitizations and exclude cross-reactivity dependent sensitization:

- Sensitization to wheat specific components supports a true food wheat allergy and help rule out clinically irrelevant sensitizations due to grass cross-reactivity.

Assess the risk for severe reactions of the wheat allergy:

- IgE antibodies to Tri a 19 and Gliadin are associated with severe reactions in wheat food allergies.

Improve patient management

- An aid to rule in IgE mediated wheat allergy as a cause of gastrointestinal symptoms.
- Improve avoidance recommendation.
- Identify patients at risk for reactions.

Make a precise assessment

ImmunoCAP Allergen Components help you differentiate between primary allergies and cross-reactivity

Make a substantiated decision

A better differentiation helps you give relevant advice and define the optimal treatment

Make a difference

More informed management helps you improve the patient's well-being and quality of life

References: 1. Keet CA et al. *The natural history of wheat allergy*. Ann Allergy Asthma Immunol. 2009 May; 102(5): 410–5. 2. Inomata N. *Wheat allergy*. Curr Opin Allergy Clin Immunol. 2009 Mar 21. 3. Palosuo K. *Update on wheat hypersensitivity*. Curr Opin Allergy Clin Immunol. 2003 Jun; 3(3): 205–9. 4. Ebisawa M et al. *Clinical utility of IgE antibodies to omega-5 gliadin in the diagnosis of wheat allergy: a pediatric multicenter challenge study*. Int Arch Allergy Immunol. 2012; 158(1): 71–6. 5. Nilsson N et al. *Combining analyses of basophil allergen threshold sensitivity, CD-sens and IgE antibodies to hydrolysed wheat, ω-5 gliadin and timothy grass enhances the prediction of wheat challenge outcome*. Int Arch Allergy Immunol. 2013 (in press). 6. Palosuo K et al. *Wheat omega-5 gliadin is a major allergen in children with immediate allergy to ingested wheat*. J Allergy Clin Immunol. 2001 Oct; 108(4): 634–8. 7. Park HJ et al. *Diagnostic Value of the Serum-Specific IgE Ratio of omega-5 Gliadin to Wheat in Adult Patients with Wheat-Induced Anaphylaxis*. Int Arch Allergy Immunol. 2012; 157(2): 147–50. 8. Pascal M et al. *Lipid transfer protein syndrome: clinical pattern, cofactor effect and profile of molecular sensitization to plant – foods and pollens*. Clin Exp Allergy. 2012 Oct; 42(10): 1529–39. 9. Daengsuwan T et al. *IgE antibodies to omega-5 gliadin in children with wheat-induced anaphylaxis*. Allergy. 2005 Apr; 60(4): 506–9. 10. Ito K et al. *IgE antibodies to omega-5 gliadin associate with immediate symptoms on oral wheat challenge in Japanese children*. Allergy. 2008 Nov; 63(11): 1536–42. 11. Kotaniemi-Syrjänen A et al. *The prognosis of wheat hypersensitivity in children*. Pediatr Allergy Immunol. 2009 Sep 30. 12. Shibata R et al. *Usefulness of specific IgE antibodies to ω-5 gliadin in the diagnosis and follow-up of Japanese children with wheat allergy*. Ann Allergy Asthma Immunol. 2011; 107(4): 337–43. 13. Morita E et al. *Food-Dependent Exercise-Induced Anaphylaxis-Importance of Omega-5 Gliadin and HMW-Glutelin as Causative Antigens for Wheat-Dependent Exercise-Induced Anaphylaxis*. Allergol Int. 2009 Oct 25; 58(4). 14. Hofmann SC et al. *IgE detection to α/β/γ-gliadin and its clinical relevance in wheat-dependent exercise-induced anaphylaxis*. Allergy. 2012; 67(11): 1457–60. 15. Tuckova L et al. *Wheat-dependent exercise-induced anaphylaxis: Pros and cons of recombinant ω-5 gliadin and glutenins, or their epitope peptides, in diagnosis*. Clinical & Experimental Allergy. 2012; 42(8): 1146–9. 16. Romano A et al. *Lipid transfer proteins: the most frequent sensitizer in Italian subjects with food-dependent exercise-induced anaphylaxis*. Clin Exp Allergy. 2012 Nov; 42(11): 1643–53. 17. Tatham AS et al. *Clin Exp Allergy*. 2008 Sep 24; 38(9): 1712–26. 18. Matricardi PM et al. *Primary versus secondary immunoglobulin E sensitization to soy and wheat in the Multi-Centre Allergy Study cohort*. Clin Exp Allergy. 2008 Mar; 38(3): 493–500. 19. Jones SM et al. *Immunologic cross-reactivity among cereal grains and grasses in children with food hypersensitivity*. J Allergy Clin Immunol. 1995 Sep; 96(3): 341–51.

thermoscientific.com/phadia

© 2013 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries. Manufacturer: Phadia AB, Uppsala Sweden.

Head office Sweden +46 18 16 50 00
Austria +43 1 270 20 20
Belgium +32 2 749 55 15
Brazil +55 11 3345 5050
China +86 21 6865 4588
Czech Republic +420 220 518 743
Denmark +45 70 23 33 06
Finland +358 9 3291 0110
France +33 1 61 37 34 30

Germany +49 761 47 8050
Hong Kong +852 2885 4613
India +91 11 4610 7555/56
Italy +39 02 64 163 411
Japan +81 3 5365 8332
Korea +82 2 2027 5400
Norway +47 21 67 32 80
Portugal +351 21 423 5350
South Africa +27 11 792 6790

52-5109-29 962-2300 1305 heart.se

Thermo
SCIENTIFIC

Part of Thermo Fisher Scientific

Thermo
SCIENTIFIC

WHEAT
Molecular Allergy



Use components to improve the diagnosis of
IgE mediated wheat food allergy

IgE mediated wheat food allergies


- Ingested wheat can cause IgE mediated wheat allergies in both children and adults.
- Immediate wheat allergy is mainly seen in children and is commonly outgrown by school-age, but remains in a subset and may cause severe reactions.¹
- In teenagers and adults, anaphylaxis may also result from ingestion of wheat in conjunction with exercise or other situations when the body is stressed (WDEIA, Wheat dependent exercise induced anaphylaxis).^{2,3}



Wheat components improve the diagnosis of wheat food allergies

Available components:

- Tri a 14, LTP
- Gliadin (α , β , γ and ω gliadins)
- Tri a 19, ω -5 gliadin




Immediate wheat allergy

Positive test results for any of the available wheat components support a diagnosis of immediate wheat allergy.

- Sensitization to Tri a 14, Tri a 19 and/or Gliadin is associated with allergic reactions to ingested wheat.⁴⁻⁸
- IgE antibodies to Tri a 19 and Gliadin are risk markers for severe reactions.^{9,10}
- Persistent IgE levels to Gliadin and Tri a 19 are associated with slower tolerance development.^{11,12}

Immediate Wheat Allergy – Suggested test profile




Gliadin gives high sensitivity for detecting wheat food allergy while Tri a 19 provides higher specificity.

Wheat dependent exercise induced anaphylaxis

WDEIA is elicited by exercise or other co-factors such as NSAID drugs, alcohol or stress after wheat intake. WDEIA patients do generally not have a history of immediate wheat allergy, and many (30 – 50 %) are also negative in extract based wheat tests. However, a majority of the WDEIA patients are sensitized to Tri a 19 and/or Gliadin.¹³⁻¹⁵

- Positive test results for Tri a 19 and/or Gliadin support a diagnosis of suspected WDEIA.¹³⁻¹⁵
- Sensitization to LTP may be associated with a risk for co-factor mediated anaphylaxis.^{8,16}



Did you know that?

- Wheat (*Triticum aestivum*) is a grass, and one of the most common causes of food allergy in children.³
- In addition to IgE mediated wheat allergies, there are other hypersensitivity disorders caused by wheat, e.g. celiac disease and irritable bowel syndrome.³
- There are many allergen components in wheat and most patients are sensitized to several of these.¹⁷
- Some grass allergic children can be misdiagnosed as wheat allergic and recommended a wheat-free diet as wheat extract tests often show up positive due to cross-reactivity between wheat and grass components e.g. profilin and CCD.¹⁸
- Grass-dependent positivity to wheat extract tests can be ruled out using specific wheat components.
- The LTP Tri a 14 lacks cross-reactivity to grass pollen allergens.
- Some patients with wheat allergy may react to other cereals such as rye and barley due to cross-reactivity between gluten proteins (gliadins and glutenins).¹⁹
- Wheat is an ingredient in processed foods other than bakery products, for instance in beer which may elicit symptoms in allergic patients.¹⁷

